

SORRY COVID

HELP PROTECT HIGH RISK PATIENTS WITH THE UPDATED SPIKEVAX® XBB.1.5 VACCINE THIS SPRING

SPIKEVAX® XBB.1.5 (andusomeran mRNA vaccine) is indicated for active immunization against coronavirus disease 2019 (COVID-19) caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) virus in individuals 6 months of age and older.¹



**STAY IN TOUCH WITH MODERNA, JOIN OUR
MAILING LIST TO RECEIVE INVITATIONS TO MEDICAL
EDUCATIONAL EVENTS, WEBINARS, AND MORE.**

Consult the Product Monograph at spikevax.com/en-CA/xb15-pm.pdf for contraindications, warnings and precautions, adverse reactions, interactions, dosing, and conditions of clinical use. The Product Monograph is also available by calling us at 1-866-MODERNA (1-866-663-3762).

References:

1. SPIKEVAX® XBB.1.5 Product Monograph. Updated 12 September, 2023. <https://spikevax.com/en-CA/xb15-pmi.pdf>
2. Guidance on an additional dose of COVID-19 vaccines in the spring. <https://www.canada.ca/en/public-health/services/publications/vaccines-immunization/national-advisory-committee-immunization-guidance-additional-dose-covid-19-vaccines-spring-2024-individuals-high-risk-severe-illness-due-covid-19.html>

© Moderna, Inc. 2024 CA-COV-2400015, 04/2024






SPIKEVAX® XBB.1.5 DOSING SCHEDULE



spikevax® XBB.1.5
andusomeran
mRNA Vaccine

Moderna's SPIKEVAX® XBB.1.5 vaccine is available in one multidose vial for administration to all individuals ages 6 months and up.¹

This season, consider reminding your patients about the importance of getting their updated COVID-19 shot. Below is the most recent dosing chart for the SPIKEVAX® XBB.1.5 vaccine to help guide you when treating your patients.¹

Age Range	COVID-19 Vaccination History	Dose Timing	Dose	Dose Volume
 12 years of age and older	Not previously vaccinated OR previously vaccinated	At least 6 months after previous COVID-19 vaccine dose	50 mcg	0.5 mL
 5 to 11 years		At least 6 months after previous COVID-19 vaccine dose	25 mcg	0.25 mL
 6 months to 4 years of age*	Not previously vaccinated	A two-dose series may be given at least 4 weeks apart	2 x 25 mcg	0.25 mL
	Previously vaccinated (1 or more previous doses)	At least 6 months after previous COVID-19 vaccine dose	25 mcg	0.25 mL

* If the child has received only 1 prior dose of a COVID-19 vaccine, SPIKEVAX® XBB.1.5 should be administered to complete the two-dose series.

** According to NACI, dose may be administered as early as 3 months.²

 **spikevax®** XBB.1.5
andusomeran
mRNA Vaccine

moderna