



# Randomized Controlled Trials & Observational Studies

Medical studies are used to explore whether an exposure or intervention **causes** a certain outcome<sup>1</sup>

A better **understanding of causation** helps **support** decision making<sup>1,2</sup>

Both **randomized controlled trials** and **observational studies** can be used to generate medical evidence<sup>2</sup>

Well-designed randomized controlled clinical trials are considered the **"gold standard"** for medical evidence<sup>1</sup>


Observational studies are a source of data for **real-world evidence (RWE)**, providing evidence in situations **outside** of carefully controlled clinical environments<sup>3</sup>



## Confounders<sup>4</sup>

Observational studies can have the issue of **confounders**

Confounders are factors associated with the outcome of interest and the other factors studied




Can lead to **non-comparable** groups and **incorrect conclusions**




A **well-designed** trial can help reduce confounders

### Example of a confounder



A well-designed randomized controlled trial helps balance confounders<sup>1,4</sup>



Adapted from Brody (2016).<sup>5</sup>

Randomization can provide a **similar distribution of confounders** across groups<sup>4</sup>

Groups should be **comparable** except for treatment<sup>1</sup>


A **treatment–outcome relationship** can often be determined<sup>4</sup>

References:  
 1. Friedman LM, Furberg CD, DeMets DL, Reboussin CBG. Fundamentals of clinical trials. 5<sup>th</sup> ed. Switzerland: Springer; 2015. 2. International Council for Harmonization of Technical Requirements for Pharmaceuticals for Human Use. Generation considerations for clinical studies (R1). Available at: [https://database.ich.org/sites/default/files/E8-R1\\_Guideline\\_Step4\\_2021\\_1006.pdf](https://database.ich.org/sites/default/files/E8-R1_Guideline_Step4_2021_1006.pdf). Accessed December 6, 2022. 3. Faries D, Zhang X, Kadziola Z, Siebert U, Kuehne F, Obenchain RL, Haro JM. Real world health care data analysis: causal methods and implementation using SAS®. Cary, NC: SAS Institute Inc.; 2020. 4. Hackshaw AK. A concise guide to observational studies in healthcare. London: Wiley; 2015. 5. Brody T. Clinical trials: study design, endpoints and biomarkers, drug safety and FDA and ICH guidelines. 2<sup>nd</sup> ed. London: Academic Press; 2016.




# Randomized Controlled Trials & Observational Studies

## Characteristics of RCTs and Observational Studies<sup>1-3</sup>




## Summary



RCTs are considered the *gold standard* for clinical research and can be used to assess efficacy of an intervention<sup>2</sup>

Observational studies *add to evidence* from RCTs – providing additional data from a *broad patient population* or subgroups of patients typically excluded from, or not represented in, RCTs<sup>1</sup>



Adapted from Faries, et al. (2020).<sup>1</sup>

References:  
 1. Faries D, Zhang X, Kadziola Z, Siebert U, Kuehne F, Obenchain RL, Haro JM. Real world health care data analysis: causal methods and implementation using SAS®. Cary, NC: SAS Institute Inc.; 2020. 2. Friedman LM, Furberg CD, DeMets DL, Reboussin CBG. Fundamentals of clinical trials. 5<sup>th</sup> ed. Switzerland: Springer; 2015. 3. Hackshaw AK. A concise guide to observational studies in healthcare. London: Wiley; 2015.

